

A Comparison of the Freshwater Macroinvertebrate Assemblages of St. Kitts and Nevis, West Indies

David Bass

Biology Department, University of Central Oklahoma, Edmond, Oklahoma, USA 73034

E-mail: dbass@ucok.edu

ABSTRACT

A survey of macroinvertebrates inhabiting the freshwater habitats of St. Kitts and Nevis was conducted during June 1996, May 1997, and January 2000. Qualitative collections were made by sweeping a dip net through the water column and by hand examination of rocks, plants, and debris submerged in both flowing and standing bodies of freshwater across these islands. These collections yielded at least 90 species, 57 from St. Kitts and 61 from Nevis. Many of the species encountered are being reported for the first time from these islands. Dominant taxa collected included several species of gastropods, decapod crustaceans, odonates, hemipterans, and coleopterans. Generally the macroinvertebrate fauna of St. Kitts and Nevis is sparse, most likely due to their oceanic origins and disturbances of their freshwater environments.

Key words: Freshwater invertebrates, St. Kitts, Nevis, Lesser Antilles.

INTRODUCTION

St. Kitts and Nevis are small Caribbean islands in the Leeward Islands of the Lesser Antilles. St. Kitts is volcanic in origin, rising 1,157 metres above sea level on Mount Liamuiga, and comprises approximately 177 km². Its southeast peninsula is much more arid than the forest-covered volcanic peaks. Aquatic habitats include shallow coastal salt ponds, small inland ponds filled with vascular plants, and mountain streams. Nevis is also volcanic in origin, rising 986 metres above sea level at Nevis Peak, and comprises approximately 94 km². This smaller island is distinguished by its central, cone-shaped volcanic peak. No permanently flowing water habitats exist in Nevis; only a few temporary streams flow on a seasonal basis. Much of the land surface is relatively porous and underground lava tubes carry water down the mountain slopes, often emerging as springs near the coasts. Standing water comprises all other freshwater habitats of Nevis. Hutton (1976) states Nevis is covered by rock laid down about two million years ago. The two islands are separated by a strait measuring less than four kilometres between the shores.

A limited amount of information regarding the freshwater invertebrates of the Lesser Antilles and other small Caribbean islands is available. Biodiversity surveys have been conducted on some islands including Barbados (Bass 2003a), St. Vincent (Harrison and Rankin 1975, 1976a, 1976b), St. Lucia (McKillop and Harrison 1980), Antigua (Bass 2005), Grenada (Flint and Sykora 1993; Bass 2004), Tobago (Nieser and Alkins-Koo 1991; Bass 2003b), and Trinidad (Hynes 1971; Alkins *et al.* 1981; Alkins-Koo 1990; Nieser and Alkins-Koo 1991). While some invertebrate groups in the Lesser Antillean region have been studied, such as decapod crustaceans (Chace and Hobbs 1969; Hart 1980), odonates (Donnelly 1970), and trichopterans (Flint 1968, 1996; Botosaneanu 2002;

Botosaneanu and Alkins-Koo 1993; Flint *et al.* 1999), many others have yet to be surveyed. A general description of freshwater habitats in Nevis and the common groups of invertebrates that inhabit them were provided by Bass (2000), but a species list was lacking and many details were omitted due to the nature of that publication. No such research from St. Kitts has been published.

The objectives of this investigation include: 1) to determine the species of aquatic macroinvertebrates inhabiting freshwater environments of St. Kitts and Nevis; 2) to note microhabitat preferences of each species; 3) to determine the relative abundance of each species; and 4) to compare the macroinvertebrates of St. Kitts and Nevis to each other and to those of other small Caribbean islands.

MATERIALS AND METHODS

Seventeen sampling sites and fourteen sampling sites were established in various freshwater habitats across St. Kitts and Nevis, respectively. These sites were visited and collections were made during June 1996, May 1997, and January 2000. Water temperature was also recorded from each site at the time of collection. Some of those sites were visited more than once.

Several methods of collecting were employed to ensure as many species as possible were captured. Submerged debris, such as stones, leaves, and wood were carefully examined and inhabitants were picked from the substrate using forceps. A dip net (mesh = 0.5 mm) was swept through aquatic vegetation and the water column to capture macroinvertebrates occupying those microhabitats. The microhabitat from where each specimen occurred was noted. Collecting efforts continued at each site until it appeared no more additional species were encountered. These collecting methods were similar to those used on

other islands so comparisons of the results could be made (Bass 2003a, 2003b, 2004, 2005). Specimen were preserved in 70% ethanol and returned to the laboratory for further identification. Taxa that could not be identified to the species level were separated into morphospecies for subsequent analysis and the taxonomic name to which they could be identified was used. Sorenson's index of similarity (1948) was used to compare these collections in St. Kitts and Nevis with similar endeavors on other small Caribbean islands.




Fig. 1. Map indicating location of collecting sites in St. Kitts and Nevis. Specific locations, dates, and approximate elevations of collections are listed in legend of Table 1.

RESULTS AND DISCUSSION

At least 90 species were recorded from these sister islands – 57 species from St. Kitts and 61 species from Nevis (Table 1). This list is important because it represents the only collections of freshwater invertebrates currently known from these two islands. Insects, especially odonates, hemipterans, and coleopterans, dominated in these collections. Gastropods were also quite common on both islands while decapod crustaceans were frequent occurrences in streams of St. Kitts. Only 27 taxa occurred on both islands.

Porifera

Radiospongilla crateriformis is the only freshwater sponge known to exist on Caribbean islands. This species was initially reported by Bass and Volkmer-Ribeiro (1998) and, in the Caribbean, has been found only on Barbados and Nevis. It was in the form of a light-colored, encrusting growth living on submerged wood debris in Hog Valley Reservoir and Old Spring Hill Pond. All of these sites from

where *R. crateriformis* has been reported in the Caribbean basin become dry during periods of extended drought and the sponge survives by forming drought-resistant gemmules until the water returns.

Platyhelminthes

Girardia was the only flatworm encountered, and it was found in only two mountain streams of St. Kitts and a temporary stream of Nevis. The only other reports of this genus in the Lesser Antilles come from Barbados (Bass 2003a) and Grenada (Bass 2004).

Oligochaeta

Only one species of oligochaete, *Allonais paraquayensis*, was collected. This worm existed in the detritus on the bottom of Hog Valley Reservoir in Nevis. This constitutes the first report of *A. paraquayensis* from a Caribbean island (M. Wetzel, personal communication).

Hirudinea

Although *Helobdella triserialis* was collected on both islands, it was limited to standing freshwater environments having dense growths of aquatic vascular plants. Based on observations by Davies (1979), it is hypothesized *H. triserialis* was passively dispersed to St. Kitts and Nevis by egg cocoons attached to macrophytes. It is also possible that aquatic birds carried ingested cocoons between different bodies of water (Davies *et al.* 1982).

Gastropoda

Nine species of freshwater snails were collected from St. Kitts and Nevis. The most abundant of these, *Physella marmorata*, occurred in ponds and streams on both islands. It is interesting that *Tropicorbis albicans* was found at several ponds having dense stands of macrophytes in Nevis, but appears to be absent in St. Kitts. Six gastropod species were collected at only one location. Notably absent from these collections is *Melanoides tuberculata*, an introduced species reported on other small islands of the eastern Caribbean basin (Bacon *et al.* 1978; Bass 2003a, 2003b, 2003c, 2004, 2005).

Pelecypoda

Eupera cubensis was found only in Hog Valley Reservoir and was one of the few species previously reported from Nevis (Bass 2000). Its distribution ranges from the southeastern United States through the Caribbean islands (Thorp and Covich 2001). This small clam is intolerant of salt water and presumably may be transported between bodies of water inadvertently attached to damp feathers of waterfowl such as herons, egrets, or coots.

Table 1. List of freshwater macroinvertebrates, including collecting sites, life cycle stages present, relative occurrence, microhabitats, and proposed trophic relationships in St. Kitts and Nevis during June 1996, May 1997, and January 2000. Life cycle: A, adult; J, juvenile; L, larva; N, nymph. Occurrence: *** abundant, ** common, * uncommon.

Taxa	Collection Sites	Life Cycle	Occurrence	Microhabitat	Trophic Relationship ¹
Porifera					
<i>Radiospongilla crateriformis</i>	N12, N20	A	*	Wood debris	Filterer
Platyhelminthes					
<i>Girardia</i> sp.	SK11, SK12, N24	A	*	Detritus	Predator
Oligochaeta					
<i>Allonais paraquayensis</i>	N12	A	*	Detritus	Detritivore
Hirudinea					
<i>Helobdella triserialis</i>	SK7, N9, N12, N23	A	**	Vascular hydrophyte	Predator
Gastropoda					
<i>Ferrissia beaufi</i>	N11	A	*	Detritus	Herbivore
<i>Helisoma</i> sp.	SK7, SK18, N15	A	*	Vascular hydrophyte	Detritivore
<i>Marisa cornuarietis</i>	SK11, SK21	A	*	Rock	Herbivore
<i>Neritina punctulata</i>	N13	A	*	Rock	Herbivore
<i>Neritina virginea</i>	SK1	A	*	Rock	Herbivore
<i>Physella marmorata</i>	SK7, SK9, SK11, SK12, SK14, SK15, SK18, SK21, N1, N5, N8, N9, N10, N14, N15, N20, N22, N23, N24	A	***	Detritus	Detritivore
<i>Planorbella (Seminolina)</i> sp.	SK7	A	*	Vascular hydrophyte	Detritivore
<i>Pyrgophorus parvulus</i>	N13	A	*	Detritus	Detritivore
<i>Tropicorbis albicans</i>	N5, N6, N8, N9, N11, N14, N19, N20, N22, N23	A	***	Detritus	Detritivore
Pelecypoda					
<i>Eupera cubensis</i>	N12, N19	A	*	Sediment	Filterer
Branchiopoda					
<i>Artemia franciscana</i>	SK3	A	*	Sediment	Filterer
Ostracoda					
<i>Chlamydotheca unispinosa</i>	N9, N12	J, A	*	Vascular hydrophyte	
Unknown Ostracoda	SK19	A	*	Detritus	
Decapoda					
<i>Atya innocous</i>	SK14, SK15, SK17, SK21	J, A	**	Detritus	Detritivore
<i>Atya</i> sp.	SK10, SK14, SK15, SK20	J	**	Detritus	Detritivore
<i>Macrobrachium crenulatum</i>	SK14, SK17, SK21	A	**	Detritus	Detritivore
<i>Macrobrachium faustinum</i>	SK10, SK17, SK20, N1, N3, N21	J	**	Detritus	Predator
<i>Micratya poeyi</i>	SK10	A	*	Detritus	Predator
<i>Palaemon pandaliformis</i>	SK1, N13, N18	J, A	*	Detritus	Detritivore
<i>Xyphocaris elongata</i>	N21	J	*	Detritus	Detritivore
Unknown Decapoda (crab)	N18	J	*	Detritus	Detritivore
Ephemeroptera					
<i>Americabaetis</i> sp.	SK10, SK11, SK12, SK14, SK15, SK17, SK21	N	***	Detritus	Collector
<i>Caenis</i> sp.	N10, N12, N15, N19	N	**	Detritus	Collector
<i>Callibaetis</i> sp.	SK7, N1, N2, N5, N6, N10, N11, N12, N24	N	***	Detritus	Collector
<i>Leptohyphes</i> sp. (2?)	SK11, SK15	N	*	Detritus	Collector
Odonata					
<i>Aeshna psilus</i>	N10	N	*	Detritus	Predator
<i>Anax concolor</i>	SK19, N13, N17, N20, N22, N23, N24	N	**	Vascular hydrophyte	Predator
<i>Brachymesia furcata</i>	SK16, N19	N	*	Detritus	Predator
<i>Brechmorhoga praecox grenadensis</i>	SK11, SK12, SK13, SK15, SK16, SK21	N	***		Predator
<i>Dythemis sterilis</i>	SK4, SK18, N10, N24	N	*	Detritus	Predator
<i>Enallagma coecum</i>	SK11, SK12	N	*	Detritus	Predator
<i>Erythemis</i> poss. <i>vesiculosa</i>	SK18, N15	N	*	Detritus	Predator
<i>Erythrodiplos</i> sp.	SK21, N2, N4, N6, N10, N24	N	***	Vascular hydrophyte	Predator
<i>Ischnura ramburii</i>	SK5, SK7, SK18, SK19, SK21, N1, N5, N9, N11, N13, N14, N15, N17, N19, N20, N21, N23	N	***	Vascular hydrophyte	Predator
<i>Lestes</i> poss. <i>forficula</i>	N11, N15, N17, N20, N23	N	***	Vascular hydrophyte	Predator
<i>Miathyria marcella</i>	SK7, SK9, N5, N15, N17	N	**	Vascular hydrophyte	Predator
<i>Orthemis ferruginea</i>	SK5, N4, N19, N20	N	**	Vascular hydrophyte	Predator

<i>Pantela flavescens</i>	N2	N	*	Detritus	Predator
<i>Tramea abdominalis</i>	SK19, N5, N10, N11, N12, N15, N20, N23	N	***	Detritus	Predator
Hemiptera					
<i>Belostoma subspinosum</i>	N5, N8	N	*	Vascular hydrophyte	Predator
<i>Buenoa</i> sp.	SK6, SK7, N6, N14	N, A	**	Water column	Predator
<i>Centrocorisa nigripennis</i> ?	N4, N10, N11, N13, N23	N, A	***	Water column	Predator
<i>Limnogonus franciscanus</i>	SK6, SK7, SK18, SK19, N13, N14, N15, N18, N21	N, A	***	Neuston	Predator
<i>Mesovelvia mulsanti</i>	SK5, SK7, N1, N2, N4, N5, N6, N8, N9, N11, N12, N13, N14, N15, N17, N18, N20, N21, N22, N23	N, A	***	Neuston	Predator
<i>Microvelia longipes</i>	N4, N7, N18	N, A	*	Neuston	Predator
<i>Microvelia</i> sp.	SK6, SK9, SK12, SK19, SK21, N1, N3, N4, N5, N7, N10, N12, N15, N16, N17, N21, N22, N24	N, A	***	Neuston	Predator
<i>Notonecta indica</i>	SK9, SK10, N2, N5, N8, N15, N19, N20, N22, N24	N, A	***	Vascular hydrophyte	Predator
<i>Pelocoris poeyi</i>	SK7, SK18, N4, N5, N1, N12, N13, N14, N15, N20, N21, N22, N23	N, A	***	Vascular hydrophyte	Predator
<i>Rhagovelia elegans</i>	SK10, SK11, SK12, SK14, SK15, SK17, SK20, SK21	N, A	***	Neuston	Predator
<i>Ranatra galatae</i>	N15, N22	A	*	Vascular hydrophyte	Predator
<i>Saldula</i> sp.	N17	N	*	Detritus	Predator
<i>Trichocorixa reticulata</i>	SK1, SK2, SK3, SK4, SK6, SK8	N, A	***	Detritus	Predator
Trichoptera					
<i>Chimarra (Chimarra)</i> sp.	SK11, SK12, SK13, SK14, SK15, N24	L	**	Detritus	Collector
<i>Smicridea (Smicridea)</i> sp.	SK12, SK14, SK15	L	**	Detritus	Collector
<i>Xiphocentron</i> sp.	SK11	L	*	Detritus	Collector
Lepidoptera					
<i>Petrophila</i> sp.	SK15	L	*	Rock	Herbivore
Coleoptera					
Aleocharinae	SK9, SK15, SK17, SK20, SK21	A	**	Detritus	Predator
<i>Copelatus postcatus</i>	SK9, SK17, N10, N22	A	**	Detritus	Predator
<i>Derallus rudis</i>	N9, N14, N23	A	**	Detritus	Collector
<i>Enochrus pseudochraceus</i>	SK10	A	*	Detritus	Herbivore
<i>Eretes sticticus</i>	SK5	A	*	Detritus	Predator
<i>Helochaeres femoratus</i>	N9, N15, N16, N17, N19, N22, N23, N24	A	***	Vascular hydrophyte	Collector
<i>Hydrobiomorpha phallica</i>	SK7	A	*	Detritus	Collector
<i>Hydrophilus insularis</i>	SK5, SK7, N2, N5, N6, N13 L,	A	***	Detritus	Predator
<i>Laccobius</i> sp.	SK18	A	*	Vascular hydrophyte	Herbivore
<i>Laccophilus subsignatus</i>	SK5, SK6, SK9, SK12, SK18, SK19, N10, N15, N19, N24	A	***	Detritus	Predator
<i>Megadytes</i> sp.	N4, N11, N20	L	**	Vascular hydrophyte	Predator
<i>Paracymus confusus</i>	N1, N4, N6, N17	A	**	Detritus	Collector
<i>Paracymus nanus</i>	SK13	A	*	Detritus	Collector
<i>Scirtes</i> sp.	N5	A	*	Vascular hydrophyte	Herbivore
<i>Thermonectes bacillaris</i>	N5, N6	L, A	*	Vascular hydrophyte	Predator
<i>Tropisternus lateralis</i>	SK7, SK18, SK19, N1, N4, N5, N6, N9, N10, N11, N12, N14, N15, N17, N20, N21, N22, N23	L, A	***	Detritus	Collector, Predator
Diptera					
<i>Ablabesmyia</i> sp.	N2, N12	L	*	Sediment	Predator
<i>Anopheles</i> sp.	N17	L	*	Water column	Collector
<i>Chironomus</i> sp.	SK5, SK16, SK19, N6, N11	L	**	Sediment	Collector
<i>Coelotanyus</i> sp.	N2, N11	L	*	Sediment	Predator
<i>Corynoneura</i> sp.	SK15	L	*	Sediment	Collector
<i>Cricotopus</i> sp.	SK12, SK15	L	*	Sediment	Collector
<i>Culex</i> sp.	N5	L	*	Water column	Collector
<i>Einfeldia</i> sp.	N12	L	*	Sediment	Collector
<i>Goeldichironomus</i> sp.	SK19	L	*	Sediment	Collector
<i>Mansonia</i> sp.	N5, N9, N14	L	**	Vascular hydrophyte	Collector
<i>Monopelopia</i> ? sp.	N11	L	*	Sediment	Predator
<i>Parachironomus</i> sp.	N11	L	*	Sediment	Collector
<i>Odontomyia</i> sp.	SK20	L	*	Detritus	Collector
<i>Paracricotopus</i> sp.	SK21	L	*	Sediment	Collector
<i>Simulium</i> sp.	SK11, SK21, N24	L	**	Rocky substrate	Collector
<i>Thienemanniella</i> sp.	SK10, SK12	L	*	Sediment	Collector

Trophic relationship¹—Trophic relationships of insects based on Merritt and Cummins (1996) and non-insects on Thorp and Covich (2001).

Collecting sites: SK1) Salt Pond, Frigate Bay, St. Kitts (2m), 25 June 1996; SK2) Salt Pond, Friars Bay Estate, St. Kitts (2m), 25 June 1996; SK3) Great Salt Pond, Salt Pond Estate (2m), 25 June 1996; SK4) Fleming Estate Pond, Fleming Estate, St. Kitts (2m), 25 June 1996; SK5) Livestock Pond, Fleming Estate, St. Kitts (4m), 25 June 1996; SK6) Muddy Pond, Frigate Bay, St. Kitts (2m), 25

June 1996; SK7) Ottley's Pond, Ottley's Estate, St. Kitts (130m), 25 June 1996; SK8) Golf Course Pond, Frigate Bay, St. Kitts (7m), 27 June 1996; SK9) Wingfield River, Wingfield Estate, St. Kitts (60m), 27 June 1996; SK10) Stone Fort River, Stone Fort Estate, St. Kitts (120m), 19 May 1997; SK11) West Farm Gut, West Farm Estate, St. Kitts (90m), 19 May 1997; SK12) Phillips River, Catchment Area, St. Kitts (290m), 20 May 1997; SK13) Ogee's River Tributary, Catchment Area, St. Kitts (310m), 20 May 1997; SK14) Frankland's River, Catchment Area, St. Kitts (380m), 22 May 1997; SK15) Wingfield River, Catchment Area, St. Kitts (155m), 22 May 1997; SK16) Dos D'ane Pond, Verchild's Mountain, St. Kitts (945m), 24 May 1997; SK17) Wingfield River, Wingfield Estate, St. Kitts (50m), 6 January 2000; SK18) Ottley's Pond, Ottley's Estate, St. Kitts (130m), 10 January 2000; SK19) Livestock Pond, Fleming Estate, St. Kitts (4m), 10 January 2000; SK20) Stone Fort River, Stone Fort Estate, St. Kitts (120m), 11 January 2000; SK21) West Farm Gut, West Farm Estate, St. Kitts (90m), 11 January 2000; N1) Golf Course Ponds, Four Seasons Resort, Nevis (7m), 26 June 1996; N2) Spring Hill Pond, Spring Hill, Nevis (170m), 26 June 1996; N3) Mount Lily River, Camps, Nevis (25m), 26 June 1996; N4) Fothergill's Estate Pond, Fothergill's Estate, Nevis (175m), 26 June 1996; N5) Pond Hill Pond, Pond Hill, Nevis (250m), 26 June 1996; N6) Nelson Spring, Cotton Ground, Nevis (2m), 26 June 1996; N7) Bath Creek, Bath, Nevis (15m), 21 May 1997; N8) Pond Hill Pond, Pond Hill, Nevis (250m), 21 May 1997; N9) Herbert Heights Pond, Herbert Heights, Nevis (360m), 21 May 1996; N10) New River Spring, New River (Road-To-Nowhere), Nevis (90m), 21 May 1997; N11) Old Spring Hill Pond, Spring Hill, Nevis (170m), 21 May 1997; N12) Hog Valley Reservoir, Hog Valley, Nevis (165m), 21 May 1997; N13) Nelson Spring Pond, Cotton Ground, Nevis (2m), 21 May 1997; N14) Pond Hill Pond, Pond Hill, Nevis (250m), 7 January 2000; N15) Fothergill's Estate Pond, Fothergill's Estate, Nevis (175m), 7 January 2000; N16) Boiling Pot, Nevis Museum, Charleston, Nevis (2m), 8 January 2000; N17) Nelson Spring, Cotton Ground, Nevis (2m), 8 January 2000; N18) Nelson Spring Pond, Cotton Ground, Nevis (2m), 8 January 2000; N19) Hog Valley Reservoir, Hog Valley, Nevis (165m), 8 January 2000; N20) Old Spring Hill Pond, Spring Hill, Nevis (170m), 8 January 2000; N21) Golf Course Ponds, Four Seasons Resort, Nevis (7m), 9 January 2000; N22) Zetlands Pond, Zetlands, Nevis (490m), 9 January 2000; N23) Herbert Heights Pond, Herbert Heights, Nevis (360m), 9 January 2000; N24) New River Spring, New River (Road-To-Nowhere), Nevis (90m), 9 January 2000.

Branchiopoda

Artemia franciscana was collected only from a warm, very shallow and somewhat isolated arm of Great Salt Pond on St. Kitts. As implied by the name of the pond, salt water was present, and the presence of *A. franciscana* in that pond indicates its ability to tolerate saline conditions. Horne and Beyenbach (1974) noted the presence of hemoglobin in *Artemia* that allows these beasts to live in aquatic habitats with low oxygen values, which probably occurred in Great Salt Pond. Birds may play a role in brachiopod dispersal as demonstrated by Proctor (1964) when *Artemia* eggs were hatched from feces of ducks.

Ostracoda

Two different ostracods were found during this investigation. *Chlamydotheca unispinosa* was collected from among submerged macrophytes at two ponds in Nevis while an unidentified ostracod was taken from a sediment sample of a pond in St. Kitts.

Decapoda

At least seven species of decapod crustaceans were encountered on these two islands. *Macrobrachium faustum* and *Palaemon pandaliformis* were the only decapods found on both islands. The atyid shrimps appear to be absent from Nevis due to the lack of running water habitats which they require. One crab was only briefly observed in Nevis, but it escaped before a specific identification could be determined. All of the species collected from St. Kitts and Nevis have been reported from other Caribbean islands (Chace and Hobbs 1969).

Ephemeroptera

A total of four mayfly genera were collected from these

two islands, although several of these occurred at more than one site. Three were limited to St. Kitts and three were found only in Nevis, while only one, *Callibaetis*, occurred on both islands. All four of these genera are widespread throughout the Lesser Antilles.

Odonata

Most of the fourteen species of odonates found were present on both St. Kitts and Nevis. Only two species and three species were limited to St. Kitts and Nevis, respectively. The most common odonate collected, the damselfly *Ischnura ramburii*, has been reported primarily in ponds on several other eastern Caribbean islands (Donnelly 1970; Harrison and Rankin, 1976b; Bass 2003a, 2003b, 2004, 2005). All of these odonates are predators that inhabit submerged vegetation and detritus.

Hemiptera

Hemipterans are one of the most frequently encountered aquatic groups of insects in the Caribbean region (Bass 2003a, 2003b, 2004, 2005). Both adults and nymphs are predators in these environments. Eight species were collected in St. Kitts and 11 species were taken from Nevis with six of those taxa being common to both islands. Some of the most common species found include the backswimmer, *Notonecta indica*, and creeping water bug, *Pelecoris poeyi*. The water boatman, *Trichocorixa reticulata*, was limited in distribution to the coastal brackish water ponds in St. Kitts where it occurred in high densities. Other frequently collected hemipterans included the water striders *Limnogonus franciscanus*, *Mesovelia mulsanti*, *Microvelia*, and *Rhagovelia elegans*. As on other small eastern Caribbean islands, these water striders had both winged and wingless adults composing the populations. This

phenomenon of wing polymorphism and its advantages to hemipterans living in isolated habitats has been well documented (Roff 1990; Schuh and Slater 1995; Thorp and Covich 2001; Bass 2003c).

Trichoptera

Trichopteran larvae are widespread across the Caribbean islands, especially in flowing water habitats (Flint 1968, 1996; Flint and Sykora 1993, Flint *et al.* 1999; Botosaneanu 2002). Although St. Kitts appears to have suitable environments to support many more species, such as in Grenada and Dominica, only three genera were encountered. It is necessary to collect adults via light-trapping, as did Flint (1968, 1996), Flint and Sykora (1993), Flint *et al.* (1999), and Botosaneanu (2002), to collect higher numbers of species, and that method was not used in this study. Therefore, it is expected more species of caddisflies will be reported from St. Kitts, and possibly Nevis, in the future. The reduced number of trichopterans on Nevis is largely due to the limited amount of flowing waters, the preferred habitat for many species of caddisfly larvae.

Lepidoptera

Larvae of *Petrophila* were collected only once. This was in the upper sections of the Wingfield River during May 1997. These larvae are widespread in mountain streams of other small eastern Caribbean islands, including Tobago (Bass 2003b) and Grenada (Bass 2004), so it is surprising this species was not collected in other streams of St. Kitts.

Coleoptera

Sixteen species of aquatic beetles were collected from St. Kitts and Nevis. Although these insects are capable of flight and presumably could exist on both islands, it is interesting to note that most of these species were found only on one island. The most commonly encountered species were the water scavenger beetle, *Tropisternus lateralis*, and the predacious diving beetle, *Laccophilus subsignatus*.

Diptera

Sixteen species of aquatic dipteran larvae were also collected from these two islands. However, nine of these taxa were encountered once with only *Chironomus* being found in more than three collections. *Chironomus* has been reported from several eastern Caribbean islands (Bass 2003a, 2003b, 2004, 2005). It is important to note that the majority of dipterans on these islands belong to a single family of midges, the Chironomidae. It is suspected this family is more common and widespread than currently reported, but midges are very small and much sediment

must be carefully examined to adequately determine their abundances and distributions. Helson *et al.* (2006) reports densities of larval chironomids exceeding 100 individuals/m² in similar streams of Trinidad.

Island biogeography theory (MacArthur and Wilson 1967) predicts St. Kitts would possess more species than Nevis because St. Kitts is a considerably larger island. However, Nevis has a slightly greater species richness. There are several possible explanations regarding this anomaly. Much of the size difference between the two islands lies with the arid southeastern peninsula of St. Kitts where little freshwater exists. If the area of the peninsula is subtracted, then the remaining mountainous region of St. Kitts is almost the same as the area of Nevis. In addition, there is more development on St. Kitts and this may have negatively impacted freshwater habitats.

Sorenson's similarity index indicates a faunal similarity of 0.46 between St. Kitts and Nevis (Tables 2-3). This is quite high, especially as compared to pair values of other small Caribbean islands (Bass 2003c). However, due to the very close proximity of these two islands, it is likely dispersal between St. Kitts and Nevis occurs in some groups so this result is possible (MacArthur and Wilson 1967). Another nearby island, Antigua, also shares somewhat higher values with St. Kitts and Nevis as well. Generally, as distance increased between the island pairs being compared, similarity values decreased. An interesting exception was Barbados which showed similarity values of 0.17 and 0.18 with St. Kitts and Nevis, respectively. All three islands have similar pond habitats and there is much overlap in species assemblages of those ponds.

Hurricane Lenny, a stochastic event, slowly moved across St. Kitts and Nevis 23-25 November, 1999. Wind

Table 2. Sorensen's index of similarity values comparing the freshwater macroinvertebrate fauna of St. Kitts to that of other small Caribbean islands, including approximate distances to those islands from St. Kitts and approximate island sizes. 0.00 = 0% common taxa and 1.00 = 100% common taxa.

Island	Approximate Distance (km)	Approximate Size (km ²)	Similarity Value
Nevis	4	94	0.46
Saba	49	13	0.02
Montserrat	75	83	0.20
Antigua	93	280	0.18
Dominica	218	751	0.13
St. Lucia	400	616	0.10
Barbados	551	430	0.17
Grenada	575	346	0.10
Tobago	700	300	0.12
Cayman Brac	1,813	37	0.00
Little Cayman	1,840	26	0.07
Grand Cayman	1,956	197	0.01
Guanaja	2,467	69	0.03

Table 3. Sorensen's index of similarity values comparing the freshwater macroinvertebrate fauna of Nevis to that of other small Caribbean islands, including approximate distances to those islands from Nevis and approximate island sizes. 0.00 = 0% common taxa and 1.00 = 100% common taxa.

Island	Approximate Distance (km)	Approximate Size (km ²)	Similarity Value
St. Kitts	4	177	0.46
Montserrat	53	83	0.11
Antigua	70	280	0.25
Saba	80	13	0.03
Dominica	200	751	0.11
St. Lucia	382	616	0.10
Barbados	538	430	0.18
Grenada	560	346	0.09
Tobago	685	300	0.10
Cayman Brac	1,844	37	0.01
Little Cayman	1,871	26	0.11
Grand Cayman	1,982	197	0.01
Guanaja	2,493	69	0.05

speeds measured approximately 215 km/hour and 30 cm of rain fell at R. L. Bradshaw Airport during that period (K. Orchard, personal communication). Large amounts of plant debris washed into streams and stream channels were scoured as a result of the heavy rainfall during that brief period. Evidence of this high discharge and a greatly reduced stream invertebrate fauna from that disturbance was observed in January 2000, six weeks after the hurricane passed. It was also observed that ponds on both islands held larger volumes of water than was present during previous visits, but their invertebrate populations appeared mostly unaffected. Hurricanes are also known to act as a mechanism to transport and re-distribute populations of invertebrates. It is suspected the occurrence of past storms may explain the presence of the sponge, *Radiospongilla crateriformis*, in Barbados and Nevis (Bass and Volkmer-Ribeiro 1998).

Both St. Kitts and Nevis are oceanic islands so their freshwater macroinvertebrate faunas had to colonize these islands from elsewhere. These immigrants must have suitable dispersal mechanisms and be able to tolerate unfavorable conditions encountered while crossing ocean waters (Bass 2003c). No endemic freshwater invertebrates that may have evolved in isolation on St. Kitts or Nevis have been discovered. However, as further investigations are conducted on these sister islands, additional species may be found and some of these might be unique to those islands.

ACKNOWLEDGEMENTS

This research began while the author was on sabbatical leave from the University of Central Oklahoma and serv-

ing as a Visiting Fulbright Professor and Research Fellow at the University of the West Indies. Partial funding for subsequent collecting trips was provided by the College of Graduate Studies and Research of the University of Central Oklahoma. Donna Bass, Courtney Bass, David Robinson (Nevis Historical and Conservation Society), and Kate Orchard (St. Christopher Historical Society) assisted with some aspects of the field work. The following individuals provided taxonomic assistance: C. Volkmer-Ribeiro (Porifera), R. Sluys (Platyhelminthes), M. J. Wetzel (Oligochaeta), D. Klemm (Hirudinea), F. G. Thompson, G.T. Watters (Mollusca), R. J. Mayer (Anostraca), C. J. Santos-Flores (Cladocera), L. D. Delorme (Ostracoda), H. H. Hobbs, III (Decapoda), M. Pescador (Ephemeroptera), R. W. Garrison (Odonata), J. T. Polhemus (Hemiptera), O. S. Flint, Jr. (Trichoptera), A. Solis (Lepidoptera), and P. J. Spangler (Coleoptera).

REFERENCES

- Alkins-Koo, M.** 1990. The aquatic fauna of two intermittent streams in the Southwestern Peninsula, Trinidad. *Living World, J. of the Trinidad and Tobago Field Naturalists' Club*, 1989/90: 36-42.
- Alkins, M., Desouza, G., Koo, M., Lip, R. L. C. and Shadid, S.** 1981. A study of the aquatic fauna of the Aripo Savannas. *Living World, J. of the Trinidad and Tobago Field Naturalists' Club*, 1981/82: 16-19.
- Bacon, P. R., Jaikaransingh, E. and Seeberan, G.** 1978. Notes on some freshwater mollusks from Nariva Swamp, Trinidad. *Living World, J. of the Trinidad and Tobago Field Naturalists' Club*, 1978: 14-15.
- Bass, D.** 2000. Freshwater invertebrates. In **D. Robinson and J. Lowery**, eds. *The Natural History of the Island of Nevis*. Ithaca, NY: Nevis Historical and Conservation Society Press. 69 p.
- Bass, D.** 2003a. Freshwater macroinvertebrates of Barbados. *J. Barbados Mus. Hist. Soc.*, 49: 269-280.
- Bass, D.** 2003b. A survey of freshwater macroinvertebrates in Tobago, West Indies. *Living World, J. Trinidad and Tobago Field Naturalists' Club*, 2003: 64-69.
- Bass, D.** 2003c. Comparison of freshwater macroinvertebrate communities from small Caribbean islands. *BioScience*, 53: 1094-1100.
- Bass, D.** 2004. A survey of freshwater macroinvertebrates on Grenada, West Indies. *Living World, J. Trinidad and Tobago Field Naturalists' Club*, 2004: 26-31.
- Bass, D.** 2005. A survey of freshwater macroinvertebrates on Antigua, West Indies. *Living World, J. Trinidad and Tobago Field Naturalists' Club*, 2005: 11-14.
- Bass, D. and Volkmer-Ribeiro, C.** 1998. *Radiospongilla crateriformis* (Porifera: Spongillidae) in the West Indies and taxonomic notes. *Iheringia, Ser. Zool.*, 85: 123-128.
- Botosaneanu, L.** 2002. An annotated checklist of caddisflies from the Caribbean islands, with distribution and bibliography (Insecta: Trichoptera). *Bulletin de la Societe Entomologique de*

France, 107 (1): 79-108.

Botosaneanu, L. and Alkins-Koo, M. 1993. The caddisflies (Insecta: Trichoptera) of Trinidad and Tobago, West Indies. *Bulletin de L'Institut Royal des Sciences Naturelles de Belgique, Entomologie*, 60: 39-48.

Chace, F. A. Jr. and Hobbs, H. H. 1969. Bredin-Archbold-Smithsonian Biological Survey of Dominica. The freshwater and terrestrial decapod crustaceans of the West Indies with special reference to Dominica. *US Natl. Mus. Bull.*, 292: 258 p.

Davies, R. W. 1979. Dispersion of freshwater leeches (Hirudinoidea) to Anticosti Island, Quebec. *Can. Field Nat.*, 93: 310-313.

Davies, R. W., Linton, L. R. and Wrona, F. J. 1982. Passive dispersal of four species of freshwater leeches by ducks. *Freshwat. Invert. Biol.*, 1: 40-44.

Donnelly, T. W. 1970. The odonata of Dominica, British West Indies. *Smithsonian Contrib. Zool.*, 37: 1-20.

Flint, O. S. Jr. 1968. Bredin-Archbold-Smithsonian Biological Survey of Dominica. 9. The Trichoptera (Caddisflies) of the Lesser Antilles. *Proc. US Natl. Mus.*, 125: 1-86.

Flint, O. S. Jr. 1996. Studies of Neotropical caddisflies LV: Trichoptera of Trinidad and Tobago. *Trans. Amer. Entomol. Soc.*, 122: 67-113.

Flint, O. S. Jr., Holzenthal, R. W., and Harris, S. C. 1999. Catalogue of the Neotropical Caddisflies (Insecta: Trichoptera). Columbus, OH: Ohio Biological Survey. 239 p.

Flint, O. S. Jr. and Sykora, J. 1993. New species and records of caddisflies (Insecta: Trichoptera) from the Lesser Antilles, with special reference to Grenada. *Annals of Carnegie Mus.*, 62: 47-62.

Harrison, A. D. and Rankin, J. J. 1975. Forest litter and stream fauna on a tropical island, St. Vincent, West Indies. *Verh. Internat. Verein. Limnol. Bd.*, 19: 1736-1745.

Harrison, A. D. and Rankin, J. J. 1976a. Hydrobiological studies of eastern Lesser Antillean islands. I. St. Vincent: freshwater habitats and water chemistry. *Arch. Hydrobiol./Suppl.*, 50: 96-144.

Harrison, A. D. and Rankin, J. J. 1976b. Hydrobiological studies of eastern Lesser Antillean islands. II. St. Vincent: freshwater fauna – its distribution, tropical river zonation and biogeography. *Arch. Hydrobiol./Suppl.*, 50: 275-311.

Hart, C. W. Jr. 1980. New freshwater shrimp records for Tobago, West Indies, with a summary of records for the Lesser

Antilles (Atyidae and Palaemonidae). *Proc. Biol. Soc. Wash.*, 93: 845-848.

Helson, J., Williams, D. D. and Turner, D. 2006. Larval chironomid community organization in four tropical rivers: human impacts and longitudinal zonation. *Hydrobiologia*, 559: 413-431.

Horne, F. R. and Beyenbach, K. W. 1974. Physiochemical features of hemoglobin of the Crustacean *Triops longicaudatus*. *Arch. Biochem. Biophys.*, 161: 369-374.

Hutton, C. O. 1976. The petrology of Nevis, Leeward Islands, West Indies. *Overseas Geology and Mineral Resources*, 52: 31 p.

Hynes, H. B. N. 1971. Zonation of the invertebrate fauna in a West Indian stream. *Hydrobiologia*, 38: 1-8.

MacArthur, R. H. and Wilson, E. O. 1967. The Theory of Island Biogeography. Princeton, NJ: Princeton University Press. 203 p.

McKillop, W. B. and Harrison, A. D. 1980. Hydrobiological studies of eastern Lesser Antillean islands. I. St. Lucia: freshwater habitats, water chemistry and distribution of freshwater mollusks. *Arch. Hydrobiol./Suppl.*, 57: 251-290.

Merritt, R. W. and Cummins, K. W. 1996. An Introduction to the Aquatic Insects of North America. Dubuque, IA: Kendall/Hunt Publ. Co. 862 p.

Nieser, N. and Alkins-Koo, M. 1991. The Water Bugs of Trinidad and Tobago. St. Augustine, Trinidad and Tobago: Zoology Department, Univ. of the West Indies, Occasional Paper No. 9.

Proctor, V. W. 1964. Viability of crustacean eggs recovered from ducks. *Ecology*, 45: 656-658.

Roff, D. A. 1990. The evolution of flightlessness in insects. *Ecol. Monogr.*, 60: 389-421.

Schuh, R. T. and Slater, J. A. 1995. True Bugs of the World (Hemiptera: Heteroptera). Classification and Natural History. Ithaca, NY: Cornell University Press. 416 p.

Sorenson, T. 1948. A method of establishing groups of equal amplitude in a plant society based on similarity of species content and its application to analysis of vegetation on Danish commons. *K. Dansk. Vidensk. Biol. Skrift*, 5: 1-34.

Thorpe, J. H. and Covich, A. P. 2001. Ecology and Classification of North American Freshwater Invertebrates. 2nd ed. San Diego, CA: Academic Press. 1056 p.